# DEPARTMENT of ENVIRONMENTAL SERVICES Water Division - Watershed Management Bureau

### LAKE TROPHIC DATA

### MORPHOMETRIC:

Lake: FISH POND	Lake Area (ha):	8.09
Town: COLUMBIA	Maximum depth (m):	2.3
County: Coos	Mean depth (m):	1.2
River Basin: Connecticut	Volume (m³):	97000
Latitude: 44°51'27" N	Relative depth:	0.7
Longitude: 71°28'52" W	Shore configuration:	1.39
Elevation (ft): 1395	Areal water load (m/yr):	29.38
Shore length (m): 1400	Flushing rate (yr <sup>-1</sup> ):	24.50
Watershed area (ha): 375.3	P retention coeff.:	0.43
<pre>% watershed ponded: 0.6</pre>	Lake type: natural	w/dam

BIOLOGICAL:	26 January 1999	4 August 1998
DOM. PHYTOPLANKTON (% TOTAL) #1	NO WINTER PLANKTON	DINOBRYON 25%
#2	ANALYZED	TABELLARIA 15%
#3		CERATIUM 15%
PHYTOPLANKTON ABUNDANCE (units/mL)		
CHLOROPHYLL-A (µg/L)		4.38
DOM. ZOOPLANKTON (% TOTAL) #1		CONOCHILUS 58%
#2		STLK PERITR CILIATE 12%
#3		ASPLACHNA 9%
ROTIFERS/LITER		408
MICROCRUSTACEA/LITER		109
ZOOPLANKTON ABUNDANCE (#/L)		602
VASCULAR PLANT ABUNDANCE		Common
SECCHI DISK TRANSPARENCY (m)		2.1 Visible on bottom
BOTTOM DISSOLVED OXYGEN (mg/L)	12.6	4.5
BACTERIA (E. coli, #/100 ml) #1		
#2		
#3		

## SUMMER THERMAL STRATIFICATION:

#### not stratified

Depth of thermocline (m): None Hypolimnion volume  $(m^3)$ : None Anoxic volume  $(m^3)$ : None

CHEMICAL:			FISH PONI	)	
	26 January 1999		4 August 1998		
DEPTH (m)	1.0		1.0		2.0
pH (units)	6.7		7.4		7.0
A.N.C. (Alkalinity)	15.4		38.9		22.8
NITRATE NITROGEN	0.29		< 0.05		< 0.05
TOTAL KJELDAHL NITROGEN	0.40		0.50		0.60
TOTAL PHOSPHORUS	0.015		0.017		0.019
CONDUCTIVITY (µmhos/cm)	55.6		87.7		87.6
APPARENT COLOR (cpu)	48		65		68
MAGNESIUM			1.72		
CALCIUM			14.6		
SODIUM			1.5		
POTASSIUM			1.16		
CHLORIDE	< 2		< 2		< 2
SULFATE	5		4		4
TN : TP	46		29		32
CALCITE SATURATION INDEX		÷	1.1		


All results in mg/L unless indicated otherwise

## TROPHIC CLASSIFICATION: 1998

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
**	2	3	1	6	Meso.

## COMMENTS:

- 1. Fish Pond was previously surveyed and classified in 1985. There was no change in trophic class and little change in trophic quality between the two dates.
- 2. Alkalinity, along with pH and calcium, were high for NH lakes. This is not a bad thing and is due to the unique geology of the area.
- 3. Easy access road with F&G boat ramp and parking located at the south end of the pond.
- 4. Seven cottages on the pond; most were set back with good buffers but one had a green manicured lawn to the water.



### FIELD DATA SHEET


LAKE: FISH POND TOWN: COLUMBIA DATE: 08/04/1998 WEATHER: SUNNY, MILD & CALM

DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION
0.1	22.2	8.4	95 %
0.5	22.0	8.4	95 %
1.0	22.5	8.2	93 %
1.5	21.5	7.6	84 %
2.0	20.8	4.5	49 %

SECCHI DISK (m): 2.1 VOB COMMENTS:
BOTTOM DEPTH (m): 2.1

TIME: 1130

\*Dissolved oxygen values are in mg/L



## AQUATIC PLANT SURVEY

LAK	E: FISH POND	TOWN: COLUMBIA	DATE: 08/04/1998	
Key	PLANT NAME		1 DUND INGE	
rey	GENERIC	COMMON	ABUNDANCE	
L	Spongilla	Freshwater sponge	Sparse	
a	Potamogeton amplifolius	Bass weed	Common	
W	Potamogeton pusillus	Pondweed	Common	
Т	Typha	Cattail	Scattered	
A	Sagittaria	Arrowhead	Sparse	
b	Scirpus	Bulrush	Sparse	
N	Nymphaea	White water lily	Sparse	
h	Hypericum	St. John's-wort	Sparse	
F	Potamogeton natans	Floating-leaf pondweed	Sparse	
g	Myrica gale	Sweet gale	Common	
Х		Sterile thread-like leaf	Sparse	
U	Utricularia	Bladderwort	Sparse	
С	Chara	Stonewort	Sparse	
t	Callitriche	Water starwort	Sparse	
S	Sparganium	Bur reed	Sparse	
f	-	Filamentous algae	Sparse	
G	Gramineae	Grass family	Sparse	
d	Dulichium arundinaceum	Three-way sedge	Sparse	
Y	Nuphar	Yellow water lily	Sparse	
		·		
OVERALL ABUNDANCE: Common				

# **GENERAL OBSERVATIONS:**

The state of the s

- 1. Both Potamogeton species were found in dense patches around the pond.
- 2. Loons, fish, heron and osprey were observed.
- 3. Wetlands were present along the inlet stream at the northern end.